

Uji Normalitas

Variables Entered/Removed^a

Model	Variables Entered	Variables Removed	Method
1	bahan-bakar, Plunger-barrel ^b	.	Enter

a. Dependent Variable: injektor

b. All requested variables entered.

Model Summary^b

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	,591 ^a	,349	,329	1,65320

a. Predictors: (Constant), bahan-bakar, Plunger-barrel

b. Dependent Variable: injektor

Coefficients^a

Model	Unstandardized Coefficients		Standardized Coefficients	t	Sig.
	B	Std. Error	Beta		
1 (Constant)	5,301	2,898		1,829	,072
Plunger-barrel	,423	,112	,421	3,786	,000
bahan-bakar	,344	,144	,265	2,384	,020

a. Dependent Variable: injektor

Residuals Statistics^a

	Minimum	Maximum	Mean	Std. Deviation	N
Predicted Value	17,7694	24,4903	21,4638	1,19297	69
Residual	-2,87650	5,53826	,00000	1,62871	69
Std. Predicted Value	-3,097	2,537	,000	1,000	69
Std. Residual	-1,740	3,350	,000	,985	69

a. Dependent Variable: injektor

NPAR TESTS

```
/K-S (NORMAL)=RES_1
/MISSING ANALYSIS
```

NPar Tests

One-Sample Kolmogorov-Smirnov Test

		Unstandardized Residual
N		69
Normal Parameters ^{a,b}	Mean	,0000000
	Std. Deviation	1,62871018
Most Extreme Differences	Absolute	,097
	Positive	,097
	Negative	-,048
Test Statistic		,097
Asymp. Sig. (2-tailed)		,183 ^c

a. Test distribution is Normal.

b. Calculated from data.

c. Lilliefors Significance Correction.